

Calabi-Yau differential equations of degree 2 and 3 and Yifan Yang's pullback

Gert Almkvist

1. Introduction.

The start of [2] was the pullback of a 5-th order differential equation, coming from diophantine approximation in number theory, to a 4-th order differential equation of Calabi-Yau type. Recently Yifan Yang has found a different way to do this pullback. His motivation came from a wish to simplify the matrices generating the monodromy group. Here we get the same pullback in a more direct way.

There are 14 hypergeometric Calabi-Yau (called CY in the sequel) differential equations (see [1],[3]), i.e. of type

$$\theta^4 + xP_1(\theta)$$

where $\theta = x \frac{d}{dx}$ and degree $P_1 = 4$. The next step is CY equations of degree 2, i.e. of type

$$\theta^4 + xP_1(\theta) + x^2P_2(\theta)$$

where degree $P_1, P_2 \leq 4$. So far we know 67 CY equations of degree 2. There are $4 \cdot 10 - 2 = 38$ Hadamard products of two second order equations of very special type (see [2] and section 3.1). We find 8 more coming from third order equations by multiplying the coefficients by $\binom{2n}{n}$. Then there are 14 Yifan Yang pullbacks of special 5-th order hypergeometric equations. Finally there are 7 sporadic equations found through a computer search by van Enkevort and van Straten (see [5]).

One can continue to ask for CY equations of degree 3. So far we know only 7 such examples. Observe that we count only equations of minimal degree under the equivalence relation having the same instanton numbers.

In section 4 we consider difference equations of order 2 and 3 (usually of degree 6), leading to CY equations after factorization in the Weyl algebra.

In the course of defining the Yifan Yang pullback we find a number of formulas for various Wronskians of the 4 solutions of a CY equation.

2. Wronskians and Yifan Yang's pullback.

2.1 The double Wronskian is almost the square.

Consider the 4-th order differential equation

$$y'''' + a_3 y''' + a_2 y'' + a_1 y' + a_0 y = 0$$

We assume it is MUM (Maximal Unipotent Monodromy) so we have the Frobenius solutions

$$\begin{aligned} y_0 &= 1 + A_1 x + \dots \\ y_1 &= y_0 \log(x) + B_1 x + \dots \\ y_2 &= \frac{1}{2} y_0 \log(x)^2 + \dots \\ y_3 &= \frac{1}{6} y_0 \log(x)^3 + \dots \end{aligned}$$

We also assume that the differential equation is Calabi-Yau (C-Y2), i.e. the coefficients satisfy

$$a_1 = \frac{1}{2} a_2 a_3 - \frac{1}{8} a_3^3 + a_2' - \frac{3}{4} a_3 a_3' - \frac{1}{2} a_3''$$

Let

$$t = \frac{y_1}{y_0}$$

Then in [1] it is proved that the condition C-Y2 is equivalent to

$$\frac{d^2}{dt^2} \left(\frac{y_2}{y_0} \right) = \frac{1}{y_0^2 \left(\frac{dt}{dx} \right)^3} \exp \left(-\frac{1}{2} \int a_3 dx \right)$$

In [2] it is proved that C-Y2 implies that

$$\begin{aligned} w_0 &= x \begin{vmatrix} y_0 & y_1 \\ y_0' & y_1' \end{vmatrix} \\ w_1 &= x \begin{vmatrix} y_0 & y_2 \\ y_0' & y_2' \end{vmatrix} \\ w_2 &= x \begin{vmatrix} y_0 & y_3 \\ y_0' & y_3' \end{vmatrix} = x \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} \\ w_3 &= \frac{x}{2} \begin{vmatrix} y_1 & y_3 \\ y_1' & y_3' \end{vmatrix} \\ w_4 &= \frac{x}{2} \begin{vmatrix} y_2 & y_3 \\ y_2' & y_3' \end{vmatrix} \end{aligned}$$

form a Frobenius basis of the solutions to a 5-th order differential equation

$$w^{(5)} + b_4 w^{(4)} + b_3 w''' + b_2 w'' + b_1 w' + b_0 w = 0$$

where

$$a_3 = \frac{2}{5}b_4 + \frac{2}{x}$$

Now consider the double Wronskian

$$u = \begin{vmatrix} w_0 & w_1 \\ w'_0 & w'_1 \end{vmatrix}$$

Differentiating $t = \frac{y_1}{y_0}$ we get

$$\frac{dt}{dx} = \frac{y_0 y'_1 - y'_0 y_1}{y_0^2} = \frac{w_0}{x y_0^2}$$

Similarly

$$\frac{d}{dt} \left(\frac{y_2}{y_0} \right) = \left(\frac{y_2}{y_0} \right)' \cdot \frac{dx}{dt} = \frac{(y_0 y'_2 - y'_0 y_2) x y_0^2}{y_0^2 w_0} = \frac{w_1}{w_0}$$

Then

$$\frac{d^2}{dt^2} \left(\frac{y_2}{y_0} \right) = \left(\frac{w_1}{w_0} \right)' \cdot \frac{dx}{dt} = \frac{x y_0^2 (w_0 w'_1 - w'_0 w_1)}{w_0^3} = \frac{x y_0^2 u}{w_0^3}$$

But by the condition equivalent to C-Y2 this is equal to

$$\frac{1}{y_0^2 \left(\frac{dt}{dx} \right)^3} \exp \left(-\frac{1}{2} \int a_3 dx \right)$$

Hence

$$u = \begin{vmatrix} w_0 & w_1 \\ w'_0 & w'_1 \end{vmatrix} = x^2 y_0^2 \exp \left(-\frac{1}{2} \int a_3 dx \right)$$

Yifan Yang has found the following pullback of the 5-th order equation

$$Y_0 = x^{5/2} u^{1/2} \exp \left(-\frac{1}{5} \int b_4 dx \right)$$

which is

$$\begin{aligned} Y_0 &= x^{5/2} x y_0 \exp \left(-\int \left(\frac{1}{4} a_3 + \frac{1}{5} b_4 \right) dx \right) \\ &= x^{7/2} y_0 \exp \left(-\int \left(\frac{1}{4} \left(\frac{2}{5} b_4 + \frac{2}{x} \right) + \frac{1}{5} b_4 \right) dx \right) \end{aligned}$$

$$= x^3 y_0 \exp\left(-\frac{3}{10} \int b_4 dx\right)$$

Hence Yifan Yang's pullback is just y_0 multiplied by a suitable factor. The remarkable fact is that for most of our 5-th order differential equations his pullback satisfies a 4-th order equation where the degree is half of the degree of the equation for y_0 . This is not the case for a general y_0 . On the contrary, the substitution

$$Y_0 = x^{3/2} y_0 \exp\left(-\frac{3}{4} \int a_3 dx\right)$$

of e.g. #209 gives an equation of degree 16 (#209 refers to the numbering in [3] also called the big table)

Example. (#130 by Helena Verrill)

Let

$$A'_n = \sum_{i+j+k+l+m+p=n} \left(\frac{n!}{i!j!k!l!m!p!} \right)^2$$

Then

$$w_0 = \sum A'_n x^n$$

satisfies the 5-th order differential equation

$$\begin{aligned} & \theta^5 - 2x(2\theta + 1)(14\theta^4 + 28\theta^3 + 28\theta^2 + 14\theta + 3) \\ & + 4x^2(\theta + 1)^3(196\theta^2 + 392\theta + 255) - 1152x^3(\theta + 1)^2(\theta + 2)^2(2\theta + 3) \end{aligned}$$

The Yifan pullback is

$$\begin{aligned} & \theta^4 - 4x \left\{ 28\left(\theta + \frac{1}{2}\right)^4 + 28\left(\theta + \frac{1}{2}\right)^2 + 1 \right\} \\ & + 3x^2 \{ 1568(\theta + 1)^4 + 2130(\theta + 1)^2 + 225 \} \\ & - 4x^3 \left\{ 23104\left(\theta + \frac{3}{2}\right)^4 + 32532\left(\theta + \frac{3}{2}\right)^2 + 3213 \right\} \\ & + x^4 \{ 872704(\theta + 2)^4 + 995680(\theta + 2)^2 + 93337 \} \\ & - 2^9 3^2 x^5 \left(\theta + \frac{5}{2}\right)^2 \left\{ 784\left(\theta + \frac{5}{2}\right)^2 + 647 \right\} \\ & + 2^{14} 3^4 x^6 (2\theta + 1)^2 \left(\theta + \frac{5}{2}\right) \left(\theta + \frac{7}{2}\right) \end{aligned}$$

which is of degree 6 while the ordinary pullback in the big table is of degree 12)

2.2 Relations between the bases of the Wronskians of the 5-th order equation and the symmetric squares of the 4-th order equation satisfying C-Y2.

The space of the Wronskians of the solutions of the 5-th order equation has dimension $\binom{5}{2} = 10$. Similarly is the dimension of the space of symmetric squares of the solutions of the 4-th order equation also $\binom{4+1}{2} = 10$ (pointed out by van Straten).

Let

$$f := \exp\left(-\frac{1}{2} \int a_3 dx\right)$$

Then we have the following explicit formulas

$$\begin{aligned} \begin{vmatrix} w_0 & w_1 \\ w'_0 & w'_1 \end{vmatrix} &= x^2 f \cdot y_0^2 \\ \begin{vmatrix} w_0 & w_2 \\ w'_0 & w'_2 \end{vmatrix} &= x^2 f \cdot y_0 y_1 \\ \begin{vmatrix} w_0 & w_3 \\ w'_0 & w'_3 \end{vmatrix} &= \frac{1}{2} x^2 f \cdot y_1^2 \\ \begin{vmatrix} w_0 & w_4 \\ w'_0 & w'_4 \end{vmatrix} &= \frac{1}{2} x^2 f \cdot (y_1 y_2 - y_0 y_3) \\ \begin{vmatrix} w_1 & w_2 \\ w'_1 & w'_2 \end{vmatrix} &= x^2 f \cdot y_0 y_2 \\ \begin{vmatrix} w_1 & w_3 \\ w'_1 & w'_3 \end{vmatrix} &= \frac{1}{2} x^2 f \cdot (y_1 y_2 + y_0 y_3) \\ \begin{vmatrix} w_1 & w_4 \\ w'_1 & w'_4 \end{vmatrix} &= \frac{1}{2} x^2 f \cdot y_2^2 \\ \begin{vmatrix} w_2 & w_3 \\ w'_2 & w'_3 \end{vmatrix} &= \frac{1}{2} x^2 f \cdot y_1 y_3 \\ \begin{vmatrix} w_2 & w_4 \\ w'_2 & w'_4 \end{vmatrix} &= \frac{1}{2} x^2 f \cdot y_2 y_3 \\ \begin{vmatrix} w_3 & w_4 \\ w'_3 & w'_4 \end{vmatrix} &= \frac{1}{4} x^2 f \cdot y_3^2 \end{aligned}$$

2.3 Higher order Wronskians.

We consider also Wronskians of order 3 and 4. Let in general

$$W(u_1, u_2, u_3) = \begin{vmatrix} u_1 & u_2 & u_3 \\ u'_1 & u'_2 & u'_3 \\ u''_1 & u''_2 & u''_3 \end{vmatrix}$$

and similarly for 4-th order Wronskians.

Proposition: We have the following identities

$$W((y_0, y_1, y_2) = f y_0$$

$$W(y_0, y_1, y_3) = f y_1$$

$$W(y_0, y_2, y_3) = f y_2$$

$$W(y_1, y_2, y_3) = f y_3$$

Proof. We have earlier proved

$$W(w_0, w_1) = x^2 f y_0^2$$

But a direct calculation shows that

$$W(w_0, w_1) = x^2 y_0 W(y_0, y_1, y_2)$$

The other identities are proved similarly.

Corollary: $W(y_0, y_1, y_2)$, etc satisfy a 4-th order differential equation with the same mirror map and Yukawa coupling as the original equation. Usually the degree of the equation is doubled.

We have the following identities

$$W(w_0, w_1, w_2) = x^3 W(y_0, y_1, y_2)^2 = x^3 f^2 y_0^2$$

$$W(w_0, w_1, w_3) = x^3 W(y_0, y_1, y_2) W(y_0, y_1, y_3) = x^3 f^2 y_0 y_1$$

$$W(w_0, w_1, w_4) = x^3 W(y_0, y_1, y_2) W(y_0, y_2, y_3) = x^3 f^2 y_0 y_2$$

$$W(w_0, w_2, w_3) = \frac{1}{2} x^3 W(y_0, y_1, y_3)^2 = x^3 f^2 y_1^2$$

$$W(w_0, w_2, w_4) = \frac{1}{2} x^3 \{W(y_0, y_1, y_2) W(y_1, y_2, y_3) + W(y_0, y_1, y_3) W(y_0, y_2, y_3)\}$$

$$= \frac{1}{2} x^3 f^2 (y_0 y_3 + y_1 y_2)$$

$$W(w_0, w_3, w_4) = \frac{1}{2} x^3 W(y_0, y_1, y_3) W(y_1, y_2, y_3) = \frac{1}{2} x^3 f^2 y_1 y_3$$

$$W(w_1, w_2, w_3) = \frac{1}{2} x^3 \{W(y_0, y_1, y_3) W(y_0, y_2, y_3) - W(y_0, y_1, y_2) W(y_1, y_2, y_3)\}$$

$$\begin{aligned}
&= \frac{1}{2}x^3 f^2(y_1 y_2 - y_0 y_3) \\
W(w_1, w_2, w_4) &= \frac{1}{2}x^3 W(y_0, y_2, y_3)^2 = \frac{1}{2}x^3 f^2 y_2^2 \\
W(w_1, w_3, w_4) &= \frac{1}{2}x^3 W(y_0, y_2, y_3)W(y_1, y_2, y_3) = \frac{1}{2}x^3 f^2 y_2 y_3 \\
W(w_2, w_3, w_4) &= \frac{1}{4}x^3 W(y_1, y_2, y_3)^2 = \frac{1}{4}x^3 f^2 y_3^2
\end{aligned}$$

We have the following formulas for 4-th order Wronskians

$$\begin{aligned}
W(w_0, w_1, w_2, w_3) &= x^3 f^3 w_0 \\
W(w_0, w_1, w_2, w_4) &= x^3 f^3 w_1 \\
W(w_0, w_1, w_3, w_4) &= x^3 f^3 w_2 \\
W(w_0, w_2, w_3, w_4) &= x^3 f^3 w_3 \\
W(w_1, w_2, w_3, w_4) &= x^3 f^3 w_4
\end{aligned}$$

We emphasize that all these formulas are only valid under the condition C-Y2.

2.4 The secret of the Yifan Yang pullback.

Consider the differential equation

$$y^{(4)} + a_3 y''' + a_2 y'' + a_1 y' + a_0 y = 0$$

which we assume is MUM and satisfying the C-Y2 condition

$$a_1 = \frac{1}{2}a_2 a_3 - \frac{1}{8}a_3^3 + a_2' - \frac{3}{4}a_3 a_3' - \frac{1}{2}a_3''$$

Then the wronskian

$$w_0 = x \begin{vmatrix} y_0 & y_1 \\ y_0' & y_1' \end{vmatrix}$$

satisfies the 5-th order equation

$$w^{(5)} + b_4 w^{(4)} + b_3 w''' + b_2 w'' + b_1 w' + b_0 w = 0$$

We have the following relations between the coefficients

$$\begin{aligned}
a_3 &= 2x^{-1} + \frac{2}{5}b_4 \\
a_2 &= -\frac{3}{2}x^{-2} + \frac{1}{2}b_3 + \frac{3}{5}x^{-1}b_4 - \frac{7}{50}b_4^2 - \frac{2}{5}b_4'
\end{aligned}$$

$$\begin{aligned}
a_1 &= \frac{3}{2}x^{-3} + b_2 + \frac{1}{2}x^{-1}b_3 - b'_3 - \frac{1}{2}b_3b_4 - \frac{3}{10}x^{-2}b_4 \\
&\quad - \frac{7}{50}x^{-1}b_4^2 + \frac{31}{250}b_4^3 - \frac{2}{5}x^{-1}b'_4 + \frac{18}{25}b_4b'_4 \\
a_0 &= -\frac{15}{16}x^{-4} - \frac{1}{4}b_1 + \frac{1}{4}x^{-1}b_2 + \frac{1}{2}b'_2 + \frac{3}{20}b_2b_4 \\
&\quad - \frac{1}{8}x^{-2}b_3 + \frac{1}{16}b_3^2 - \frac{1}{8}x^{-1}b'_3 - \frac{3}{8}b''_3 - \frac{1}{10}x^{-1}b_3b_4 \\
&\quad - \frac{21}{200}b_3b_4^2 - \frac{7}{20}b_3b'_3 - \frac{13}{40}b'_3b_4 + \frac{3}{20}x^{-3}b_4 + \frac{7}{200}x^{-2}b_4^2 \\
&\quad + \frac{11}{500}x^{-1}b_4^3 + \frac{221}{10000}b_4^4 + \frac{1}{10}x^{-2}b'_4 - \frac{1}{20}x^{-1}b''_4 + \frac{1}{10}b'''_4 \\
&\quad + \frac{3}{50}x^{-1}b_4b'_4 + \frac{29}{100}b_4b''_4 + \frac{33}{100}(b'_4)^2 + \frac{69}{250}b_4^2b'_4
\end{aligned}$$

We express the condition C-Y2 in the coefficients b_2, b_3, b_4

$$U \equiv -b_2 + \frac{3}{2}b'_3 + \frac{3}{5}b_3b_4 - b''_4 - \frac{6}{5}b_4b'_4 - \frac{4}{25}b_4^3 = 0$$

Now we assume the 5-th order equation given satisfying the condition just stated and we want to find the 4-th order equation somewhat modified to get lower degrees of the coefficients. To achieve this we will make the transformation

$$u = f \cdot y$$

Then the 4-th order equation is transformed to

$$u^{(4)} + c_3u''' + c_2u'' + c_1u' + c_0u = 0$$

where

$$\begin{aligned}
c_3 &= 4\frac{f'}{f} + a_3 \\
c_2 &= 6\frac{f''}{f} + 3a_3\frac{f'}{f} + a_2 \\
c_1 &= 4\frac{f'''}{f} + 3a_3\frac{f''}{f} + 2a_2\frac{f'}{f} + a_1 \\
c_0 &= \frac{f^{(4)}}{f} + a_3\frac{f'''}{f} + a_2\frac{f''}{f} + a_1\frac{f'}{f} + a_0
\end{aligned}$$

To simplify

$$a_3 = \frac{2}{x} + \frac{2}{5}b_4$$

we choose

$$f = x^{-1/2} \exp(\lambda \int b_4 dx)$$

with

$$\begin{aligned} \frac{f'}{f} &= -\frac{1}{2x} + \lambda b_4 \\ \frac{f''}{f} &= \frac{3}{4}x^{-2} - \lambda x^{-1}b_4 + \lambda^2 b_4^2 + \lambda b_4' \\ \frac{f'''}{f} &= -\frac{15}{8}x^{-3} + \frac{9}{4}\lambda x^{-2}b_4 - \frac{3}{2}\lambda^2 x^{-1}b_4^2 + \lambda^3 b_4^3 - \frac{3}{2}\lambda x^{-1}b_4' + \lambda b_4'' \\ \frac{f^{(4)}}{f} &= \frac{105}{16}x^{-4} - \frac{15}{2}\lambda x^{-3}b_4 + \frac{9}{2}\lambda^2 x^{-2}b_4^2 - 2\lambda^3 x^{-1}b_4^3 + \lambda^4 b_4^4 \\ &\quad + \frac{9}{2}\lambda x^{-2}b_4' - 2\lambda x^{-1}b_4'' + \lambda b_4''' + 4\lambda^2 b_4 b_4'' + 6\lambda^3 b_4^2 b_4' + 3\lambda^2 (b_4')^2 \end{aligned}$$

We get rid of the term $\frac{2}{x}$ in a_3

$$\begin{aligned} c_3 &= \frac{2(10\lambda + 1)}{5}b_4 \\ c_2 &= \frac{1}{2}b_3 + (6\lambda^2 + \frac{6}{5}\lambda - \frac{7}{50})b_4^2 + (6\lambda - \frac{2}{5})b_4' \end{aligned}$$

We want to avoid higher powers and derivatives of b_4 (remember that the b_k are rational functions so differentiating also increases the degree). We can also add multiples of the relation U (and its derivatives). Hence consider

$$\begin{aligned} c_1 + \mu U &= \\ (1 - \mu)b_2 + (-1 + \frac{3}{2}\mu)b_3' + (\lambda + \frac{3}{5}\mu - \frac{1}{2})b_3b_4 \\ &\quad + (4\lambda^3 + \frac{6}{5}\lambda^2 - \frac{7}{5}\lambda + \frac{31}{250} - \frac{4}{25}\mu)b_4^3 \\ &\quad + (12\lambda^2 + \frac{2}{5}\lambda + \frac{18}{25} - \frac{6}{5}\mu)b_4b_4' \\ &\quad + (4\lambda - \mu + \frac{2}{5})b_4'' \end{aligned}$$

By choosing λ and μ such that the three last terms vanish we get a very simple formula for c_1 agreeing with Yifan's formula

$$4\lambda^3 + \frac{6}{5}\lambda^2 - \frac{7}{5}\lambda + \frac{31}{250} - \frac{4}{25}\mu = 0$$

$$12\lambda^2 + \frac{2}{5}\lambda + \frac{18}{25} - \frac{6}{5}\mu = 0$$

$$4\lambda - \mu + \frac{2}{5} = 0$$

with the solution

$$\lambda = \frac{3}{10}$$

$$\mu = \frac{8}{5}$$

giving

$$c_1 = -\frac{3}{5}b_2 + \frac{7}{5}b'_3 + \frac{19}{25}b_3b_4$$

Fortunately $\lambda = \frac{3}{10}$ works also for c_0 . Indeed

$$c_0 + \frac{2}{5}U' - \frac{1}{4x}U + \frac{41}{100}b_4U =$$

$$-\frac{1}{4}b_1 + \frac{1}{10}b'_2 + \frac{1}{25}b_2b_4 + \frac{9}{40}b''_3 + \frac{1}{16}b_3^2 + \frac{1}{25}b_3b'_4$$

$$+ \frac{23}{100}b'_3b_4 + \frac{9}{250}b_3b_4^2$$

which agrees with Yifan Yang's formula. To get the analytic solution we have to multiply the solution by $x^{5/2}$ which is the same thing as the substitution $\theta \longrightarrow \theta - \frac{5}{2}$ in the differential equation.

Definition, The *Yifan Yang pullback* of the 5-th order equation

$$y^{(5)} + b_4y^{(4)} + b_3y''' + b_2y'' + b_1y' + b_0y = 0$$

is

$$y^{(4)} + c_3y''' + c_2y'' + c_1y' + c_0y = 0$$

where

$$c_3 = \frac{8}{5}b_4$$

$$c_2 = \frac{1}{2}b_3 + \frac{7}{5}b'_4 + \frac{19}{25}b_4^2$$

$$c_1 = -\frac{3}{5}b_2 + \frac{7}{5}b'_3 + \frac{19}{25}b_3b_4$$

$$c_0 = -\frac{1}{4}b_1 + \frac{1}{10}b'_2 + \frac{1}{25}b_2b_4 + \frac{9}{40}b''_3 + \frac{1}{16}b_3^2$$

$$+\frac{1}{25}b_3b_4+\frac{23}{100}b'_3b_4+\frac{9}{250}b_3b_4^2$$

3. Calabi-Yau differential equations of degree 2 and 3.

In the big table [3] of CY equations there are now 336 entries. Except for the first 16 there is no order in the table. The entries were put in as they were found. Here we will start to bring some order into the chaos.

Given a (MUM) differential operator L in standard form

$$L = \theta^k + xP_1(\theta) + \dots + x^dP_d(\theta)$$

where $\deg(P_j) \leq k$ we call k the *order* and d the *degree* of L . For a CY equation of degree d

$$L = \theta^4 + xP_1(\theta) + \dots + x^dP_d(\theta)$$

having Frobenius solutions y_0, y_1, y_2, y_3 we define

$$q = \exp(y_1/y_0)$$

$$K(q) = (q \frac{d}{dq})^2(y_2/y_0) = 1 + \sum_{k=1}^{\infty} \frac{k^3 N_k q^k}{1 - q^k}$$

the *Yukawa coupling*. The N_k are called *instanton numbers*. One of the conditions for being a CY equation is that there should exist a fix integer c such that cN_k are integers for all k . It is clear that q and K are invariant under the transformation

$$y(x) \rightarrow f(x)y(x)$$

One can verify that K (but not q) is also invariant under

$$y(x) \rightarrow y(g(x))$$

where

$$g(x) = x + c_2x^2 + c_3x^3 + \dots$$

We call two CY differential equations *equivalent* ($L_1 \sim L_2$) if they have the same $K(q)$. So $K(q)$ is the invariant we shall use to classify CY equations. This is also done in the "Superseeker" in the big table, where the equations are ordered after the absolute values of N_1 and N_3 (observe that the instanton numbers are not invariant under the transformation $x \rightarrow cx$ if $c \neq 1$). There are many equivalent equations in the table. Now we will only give the representative of lowest degree. Thus #129 in the table is of degree 24 but it is equivalent to #8, which in its turn will be shown to be equivalent to $\tilde{8}$, a Yifan Yang pullback of degree 2.

There are equivalences between various Hadamard products, for some of which we can even find explicit transformations. This will be treated at the end of this section.

3.1 Hadamard products.

We start with second order equations. They are of two types

(i) Hypergeometric.

They are of type

$$\theta^2 - xQ(\theta)$$

where $Q(\theta)$ is a polynomial of degree 2

Name	$Q(\theta)$
A	$4(2\theta + 1)^2$
B	$3(3\theta + 1)(3\theta + 2)$
C	$4(4\theta + 1)(4\theta + 3)$
D	$12(6\theta + 1)(6\theta + 5)$

(ii) Second degree

They are of type

$$\theta^2 - xP(\theta) - cx^2(\theta + 1)^2$$

where

$$P(\theta) = a\theta^2 + a\theta + b$$

Name	a	b	c
(a)	7	2	8
(b)	11	3	1
(c)	10	3	-9
(d)	12	4	-32
(e)	32	12	-256
(f)	9	3	-27
(g)	17	6	-72
(h)	54	21	-729
(i)	128	52	-4096
(j)	864	372	-186624

Most of these differential equations appear in Zagier's secret paper [7] and the rest were found by van Enkevort and Van Straten in a computer search. More information can be found in [2] where formulas for the coefficients are given. The Hadamard product is defined by

$$\sum A_n x^n * \sum B_n x^n = \sum A_n B_n x^n$$

We use $L_1 * L_2$ for the corresponding differential operator annihilating $\sum A_n B_n x^n$. Then we have the formula

$$(\theta^2 - xP(\theta) - cx^2(\theta + 1)^2) * (\theta^2 - xQ(\theta))$$

$$= \theta^4 - xP(\theta)Q(\theta) - cx^2Q(\theta)Q(\theta + 1)$$

*	A	B	C	D
a	45	15	68	62
b	25	24	51	63
c	58	70	69	64
d	36	48	38	65
e	111=10** ~10*	110	30~3	112
f	133	134	135	136
g	137	138	139	140
h	141=8**=8*	142	K(q)=1	143
i	7** ~7*	B*(i)	C*(i)~6*	D*(i)
j	9** ~9*	B*(j)	C*(j)	D*(j)

We get 40 degree 2 CY equations $(A) * (a), (A) * (b), \dots, (D) * (j)$ but two of them, are deleted, since $(C) * (h)$ has trivial $K(q)$. and $C * e \sim 3$. We compute the last equation.

Example. $(D) * (j)$

We have for (D)

$$Q(\theta) = 12(6\theta + 1)(6\theta + 5)$$

where

$$\theta^2 - xQ(\theta)$$

has solution

$$y_0 = \sum \frac{(6n)!}{(3n)!(2n)!n!} x^n$$

We have for (j)

$$\theta^2 - 12x(72\theta^2 + 72\theta + 31) + 186624x^2(\theta + 1)^2$$

with solution

$$y_0 = \sum_{n=0}^{\infty} 432^n \sum_{k=0}^n (-1)^k \binom{-5/6}{k} \binom{-1/6}{n-k}^2 x^n$$

We get the Hadamard product $(D) * (j)$

$$\begin{aligned} &\theta^4 - 144x(6\theta + 1)(6\theta + 5)(72\theta^2 + 72\theta + 31) \\ &+ 12^2 186624x^2(6\theta + 1)(6\theta + 5)(6\theta + 7)(6\theta + 11) \end{aligned}$$

with the solution

$$y_0 = \sum A_n x^n$$

where

$$A_n = \frac{(6n)!}{((3n)!(2n)!n!)} \sum_{k=0}^n (-1)^k \binom{-5/6}{k} \binom{-1/6}{n-k}^2$$

There are also 10 third order equations $\alpha, \beta, \dots, \varkappa$. (see [2]). Multiplying the coefficients by $\binom{2n}{n}$ we get the following 4-th order equations

$$\theta^4 - x(2\theta + 1)^2 P(\theta) + cx^2(\theta + 1)^2(2\theta + 1)(2\theta + 3)$$

where the polynomial $P(n)$ and c are given in the following table

Name	#	P(n)	c
α	16	$4(5n^2 + 5n + 2)$	256
γ	29	$16(2n^2 + 2n + 1)$	1024
δ	41	$2(7n^2 + 7n + 3)$	324
ε	42	$8(3n^2 + 3n + 1)$	64
ζ	185	$6(3n^2 + 3n + 1)$	-108
η	184	$2(11n^2 + 11n + 5)$	500
ι	4*	$6(9n^2 + 9n + 5)$	2916
$\varkappa$	13*	$48(18n^2 + 18n + 13)$	746496

We delete $\binom{2n}{n} * (\beta) \sim 3$ and $\binom{2n}{n} * (\vartheta) = 10^* \sim A * (e)$.

3.2 Yifan Yang pullbacks of hypergeometric 5-th order differential equations.

Consider the 5-th order hypergeometric differential equation

$$L_5 = \theta^5 - cx(\theta + a_1)(\theta + a_2)(\theta + a_3)(\theta + a_4)(\theta + a_5)$$

Proposition 1.

If

$$a_1 = \frac{1}{2}$$

$$a_3 = 1 - a_2$$

$$a_5 = 1 - a_4$$

then L_5 satisfies the C-Y condition

Proof: Write L_5 as

$$w^{(5)} + b_4 w^{(4)} + b_3 w^{(3)} + b_2 w'' + b_1 w' + b_0 w = 0$$

where

$$b_4 = \frac{5}{2} \cdot \frac{4 - 5cx}{x(1 - cx)}$$

$$b_3 = \frac{25 - c(42 + a_2 + a_4 - a_2^2 - a_4^2)x}{x^2(1 - cx)}$$

$$b_2 = \frac{3}{2} \cdot \frac{10 - c(26 + 3a_2 + 3a_4 - 3a_2^2 - 3a_4^2)x}{x^3(1 - cx)}$$

satisfy C-Y, i.e.

$$b_2 = \frac{3}{2}b'_3 + \frac{3}{5}b_3b_4 - b''_4 - \frac{6}{5}b_4b'_4 - \frac{4}{25}b_4^3$$

Proposition 2.

L_5 has the solutions

$$w_0 = \sum_{n=0}^{\infty} \frac{(a_1)_n (a_2)_n (a_3)_n (a_4)_n (a_5)_n}{n!^5} (cx)^n$$

$$w_1 = w_0 \log(x)$$

$$+ \sum_{n=1}^{\infty} \frac{(cx)^n}{n!^5} \prod_{j=1}^5 (a_j)_n \left\{ -5H_n + \sum_{j=1}^5 (\psi(n + a_j) - \psi(a_j)) \right\}$$

where

$$H_n = \sum_{j=1}^n \frac{1}{j}$$

$$\psi(a) = \frac{\Gamma'(a)}{\Gamma(a)}$$

Observe that

$$\psi(n + a) - \psi(a) = \sum_{j=0}^{n-1} \frac{1}{a + j}$$

$$(a)'_n = (a)_n (\psi(n + a) - \psi(a)) = \sum_{j=0}^{n-1} a(a+1) \dots \widehat{(a+j)} \dots (a+n-1)$$

Using Yinfan Yang's definition of the pullback we get the following result

Theorem

The differential equation

$$\begin{aligned} & \theta^4 - cx \left\{ 2\left(\theta + \frac{1}{2}\right)^4 + \frac{1}{2}\left(\frac{7}{2} - \alpha^2 - \beta^2\right)\left(\theta + \frac{1}{2}\right)^2 + \frac{1}{16} - \frac{1}{4}\left(\alpha^2 + \frac{1}{4}\right)\left(\beta^2 + \frac{1}{4}\right) \right\} \\ & + c^2 x^2 \left(\theta + 1 + \frac{\alpha + \beta}{2}\right) \left(\theta + 1 - \frac{\alpha + \beta}{2}\right) \left(\theta + 1 + \frac{\alpha - \beta}{2}\right) \left(\theta + 1 - \frac{\alpha - \beta}{2}\right) \end{aligned}$$

has the solution

$$y_0 = \frac{1}{\sqrt{1 - cx}} \left\{ \frac{\sum_{n=0}^{\infty} (cx)^n \sum_{k=0}^n \frac{\prod_{j=1}^5 (a_j)_k (a_j)_{n-k}}{k!^5 (n-k)!^5}}{[1 + (2k - n) \{-5H_k + \sum_{j=1}^5 (\psi(k + a_j) - \psi(a_j))\}]} \right\}^{1/2}$$

where

$$\begin{aligned}
a_1 &= \frac{1}{2} \\
a_2 &= \frac{1}{2} + \alpha \\
a_3 &= \frac{1}{2} - \alpha \\
a_4 &= \frac{1}{2} + \beta \\
a_5 &= \frac{1}{2} - \beta
\end{aligned}$$

We get the following 14 cases giving CY equations

Name	a_2	a_4	c	α	β
$\widetilde{1}$	1/5	2/5	$4 \cdot 5^5$	-3/10	-1/10
$\widetilde{2}$	1/10	3/10	$4 \cdot 8 \cdot 10^5$	-2/5	-1/5
$\widetilde{3}$	1/2	1/2	$4 \cdot 256$	0	0
$\widetilde{4}$	1/3	1/3	$4 \cdot 3^6$	-1/6	-1/6
$\widetilde{5}$	1/2	1/3	$4 \cdot 432$	0	-1/6
$\widetilde{6}$	1/2	1/4	$4 \cdot 2^{10}$	0	-1/4
$\widetilde{7}$	1/8	3/8	$4 \cdot 2^{16}$	-3/8	-1/8
$\widetilde{8}$	1/6	1/3	$4 \cdot 11664$	-1/3	-1/6
$\widetilde{9}$	1/12	5/12	$4 \cdot 12^6$	-5/12	-1/12
$\widetilde{10}$	1/4	1/4	$4 \cdot 2^{12}$	-1/4	-1/4
$\widetilde{11}$	1/4	1/3	$4 \cdot 12^3$	-1/4	-1/6
$\widetilde{12}$	1/6	1/4	$4 \cdot 2^{10} 3^3$	-1/3	-1/4
$\widetilde{13}$	1/6	1/6	$4 \cdot 2^8 3^6$	-1/3	-1/3
$\widetilde{14}$	1/2	1/6	$4 \cdot 2^8 3^3$	0	-1/3

We have the following table of the differential equations

$\widetilde{1}$

$$\begin{aligned}
&\theta^4 - 10x(2500\theta^4 + 5000\theta^3 + 5875\theta^2 + 3375\theta + 738) \\
&+ 62500z^2(5\theta + 4)(5\theta + 6)(10\theta + 9)(10\theta + 11)
\end{aligned}$$

$\widetilde{2}$

$$\begin{aligned}
&\theta^4 - 8x(80000\theta^4 + 160000\theta^3 + 186000\theta^2 + 106000\theta + 22811) \\
&+ 2^{10}10^6z^2(10\theta + 7)(10\theta + 9)(10\theta + 11)(10\theta + 13)
\end{aligned}$$

$\widetilde{3}$

$$\theta^4 - 16x(128\theta^4 + 256\theta^3 + 304\theta^2 + 176\theta + 39) + 2^{20}x^2(\theta + 1)^4$$

$\widetilde{4}$

$$\begin{aligned} &\theta^4 - 18x(324\theta^4 + 648\theta^3 + 765\theta^2 + 441\theta + 97) \\ &\quad + 2^2 3^{10} x^2 (\theta + 1)^2 (6\theta + 5)(6\theta + 7) \end{aligned}$$

 $\widetilde{5}$

$$\begin{aligned} &\theta^4 - 12x(288\theta^4 + 576\theta^3 + 682\theta^2 + 394\theta + 87) \\ &\quad + 144x^2(12\theta + 11)^2(12\theta + 13)^2 \end{aligned}$$

 $\widetilde{6}$

$$\begin{aligned} &\theta^4 - 16x(512\theta^4 + 1024\theta^3 + 1208\theta^2 + 696\theta + 153) \\ &\quad + 4096x^2(8\theta + 7)^2(8\theta + 9)^2 \end{aligned}$$

 $\widetilde{7}$

$$\begin{aligned} &\theta^4 - 16x(32768\theta^4 + 65536\theta^3 + 76544\theta^2 + 43776\theta + 9495) \\ &\quad + 2^{26} z^2 (4\theta + 3)(4\theta + 5)(8\theta + 7)(8\theta + 9) \end{aligned}$$

 $\widetilde{8}$

$$\begin{aligned} &\theta^4 - 36x(2592\theta^4 + 5184\theta^3 + 6066\theta^2 + 3474\theta + 755) \\ &\quad + 2^4 3^{10} z^2 (4\theta + 3)(4\theta + 5)(12\theta + 11)(12\theta + 13) \end{aligned}$$

 $\widetilde{9}$

$$\begin{aligned} &\theta^4 - 144x(165888\theta^4 + 331776\theta^3 + 386496\theta^2 + 220608\theta + 47711) \\ &\quad + 2^{22} 3^{10} z^2 (4\theta + 3)(4\theta + 5)(6\theta + 5)(6\theta + 7) \end{aligned}$$

 $\widetilde{10}$

$$\begin{aligned} &\theta^4 - 16x(2048\theta^4 + 4096\theta^3 + 4800\theta^2 + 2752\theta + 599) \\ &\quad + 2^{24} z^2 (\theta + 1)^2 (4\theta + 3)(4\theta + 5) \end{aligned}$$

 $\widetilde{11}$

$$\begin{aligned} &\theta^4 - 12x(1152\theta^4 + 2304\theta^3 + 2710\theta^2 + 1558\theta + 341) \\ &\quad + 144z^2(24\theta + 19)(24\theta + 23)(24\theta + 25)(24\theta + 29) \end{aligned}$$

 $\widetilde{12}$

$$\begin{aligned} &\theta^4 - 48x(4608\theta^4 + 9216\theta^3 + 10744\theta^2 + 6136\theta + 1325) \\ &\quad + 2^{12} 3^2 z^2 (24\theta + 17)(24\theta + 23)(24\theta + 25)(24\theta + 31) \end{aligned}$$

 $\widetilde{13}$

$$\begin{aligned} &\theta^4 - 12^2 x(10368\theta^4 + 20736\theta^3 + 24048\theta^2 + 13680\theta + 2927) \\ &\quad + 12^{10} z^2 (\theta + 1)^2 (3\theta + 2)(3\theta + 4) \end{aligned}$$

 $\widetilde{14}$

$$\begin{aligned} &\theta^4 - 48x(1152\theta^4 + 2304\theta^3 + 2704\theta^2 + 1552\theta + 339) \\ &\quad + 2^{16} 3^2 z^2 (6\theta + 5)^2 (6\theta + 7)^2 \end{aligned}$$

We have the equivalences $\widehat{m} \sim \widetilde{m}$ for $m=1,2,\dots,14$. Some of the Hadamard products are also equivalent to certain $\widetilde{m}$, due to the following table θ

*	(e)	(h)	(i)	(j)
(e)	$\widetilde{3}$	$\widetilde{5}$	$\widetilde{6}$	$\widetilde{14}$
(h)		$\widetilde{4}$	$\widetilde{11}$	$\widetilde{8}$
(i)			10	12
(j)				$\widetilde{13}$

Let $y_0(x)$ be the solution of $(e) * (e)$ and $\widetilde{y}_0(x)$ of $\widetilde{3}$. Then we have the transformation

$$\widetilde{y}_0(x) = f(x)y_0(g(x))$$

where

$$f(x) = 1 + 480x + 383488x^2 + 330493952x^3 + \dots$$

$$g(x) = x + 128x^2 + 91920x^3 + 52555776x^4 + \dots$$

We have not found any explicit formulas for f and g . On the contrary, if $\widehat{y}_0(x)$ is the solution of $\widehat{3}$ then

$$y_0(x) = \frac{1}{\sqrt{1 - 2^{16}x^2}} \widehat{y}_0\left(\frac{x}{(1 + 2^8x)^2}\right)$$

We also notice the following equivalences

*	β	ι	θ	$\varkappa$
A	$A^*(e)$	$B^*(e)$	$C^*(e)$	$D^*(e)$
B	$A^*(h)$	$B^*(h)$	$C^*(h)$	$D^*(h)$
C	$A^*(i)$	$B^*(i)$	$C^*(i)$	$D^*(i)$
D	$A^*(j)$	$B^*(j)$	$C^*(j)$	$D^*(j)$

Here $\beta, \iota, \theta, \varkappa$ refer to third degree differential equations in [2] and e.g. the upper most left entry means that the pullback of the 5-th order equation $A * \beta$ is equivalent to the 4-th order, degree 2 equation $A * (e)$

Hence we have replaced $14+10+16=40$ higher degree equations in the table with equations of degree 2.

3.3. Sporadic CY equations of degree 2.

The following CY equations were found in a computer search by van Enckevort and van Straten (see [3] and [5]). They found also $\#181=\widetilde{4}$, $\#204=\widetilde{3}$ and $\#31\sim 3$.

18.

$$\theta^4 - 4x(2\theta + 1)^2(3\theta^2 + 3\theta + 1) - 16x^2(2\theta + 1)(2\theta + 3)(4\theta + 3)(4\theta + 5)$$

26.

$$\theta^4 - 2x(2\theta + 1)^2(13\theta^2 + 13\theta + 4) - 12x^2(2\theta + 1)(2\theta + 3)(3\theta + 2)(3\theta + 4)$$

28.

$$\theta^4 - x(65\theta^4 + 130\theta^3 + 105\theta^2 + 40\theta + 6) + 4x^2(\theta + 1)^2(4\theta + 3)(4\theta + 5)$$

84.

$$\theta^4 - 4x(32\theta^4 + 64\theta^3 + 63\theta^2 + 31\theta + 6) + 256x^2(\theta + 1)^2(4\theta + 3)(4\theta + 5)$$

182.

$$\theta^4 - x(43\theta^4 + 86\theta^3 + 77\theta^2 + 34\theta + 6) + 12x^2(\theta + 1)^2(6\theta + 5)(6\theta + 7)$$

183.

$$\theta^4 - 4x(2\theta + 1)^2(7\theta^2 + 7\theta + 3) + 48x^2(2\theta + 1)(2\theta + 3)(4\theta + 3)(4\theta + 5)$$

205.

$$\theta^4 - x(59\theta^4 + 118\theta^3 + 105\theta^2 + 46\theta + 8) + 96x^2(\theta + 1)^2(3\theta + 2)(3\theta + 4)$$

We deleted #31 which is equivalent to #3 which is of degree one.

There are a number of 4-th order differential equations that fulfill all the conditions of CY equations but they are factorable. Most of them were found by van Enkevort and van Straten. We list them here hoping that somebody can give a geometric explanation of their existence.

(i)

$$\theta^4 - 6x(36\theta^4 + 72\theta^3 + 83\theta^2 + 47\theta + 10) + 36x^2(3\theta + 2)(3\theta + 4)(6\theta + 5)(6\theta + 7)$$

(ii)

$$\theta^4 - 4x(32\theta^4 + 64\theta^3 + 74\theta^2 + 42\theta + 9) + 16x^2(4\theta + 3)^2(4\theta + 5)^2$$

(iii)

$$\theta^4 - 4x(128\theta^4 + 256\theta^3 + 294\theta^2 + 166\theta + 35) + 16x^2(8\theta + 5)(8\theta + 7)(8\theta + 9)(8\theta + 11)$$

(iv)

$$\theta^4 - 12x(6\theta + 1)(6\theta + 5)(2\theta^2 + 2\theta + 1) + 144x^2(6\theta + 1)(6\theta + 5)(6\theta + 7)(6\theta + 11)$$

(v)

$$\theta^4 - 3x(3\theta + 1)(3\theta + 2)(2\theta^2 + 2\theta + 1) + 9x^2(3\theta + 1)(3\theta + 2)(3\theta + 4)(3\theta + 5)$$

(vi)

$$\theta^4 - 4x(4\theta + 1)(4\theta + 3)(2\theta^2 + 2\theta + 1) + 16x^2(4\theta + 1)(4\theta + 3)(4\theta + 5)(4\theta + 7)$$

(vii)

$$\theta^4 - 4x(2\theta + 1)^2(2\theta^2 + 2\theta + 1) + 16x^2(2\theta + 1)^2(2\theta + 3)^2$$

(viii)

$$\theta^4 - 2x(2\theta + 1)^2(\theta^2 + \theta + 1) + 4x^2(\theta + 1)^2(2\theta + 1)(2\theta + 3)$$

3.4. CY equations of degree 3.

So far we have only found the following CY equations of degree 3. The notation $\widetilde{214}$ means an equation which is equivalent to 214.

34.(Verrill)

$$\begin{aligned} &\theta^4 - x(35\theta^4 + 70\theta^3 + 63\theta^2 + 28\theta + 5) \\ &+ x^2(\theta + 1)^2(259\theta^2 + 518\theta + 285) - 225x^3(\theta + 1)^2(\theta + 2)^2 \end{aligned}$$

$$A_n = \sum_{i+j+k+l+m=n} \left(\frac{n!}{i!j!k!l!m!} \right)^2$$

145.

$$\begin{aligned} & \theta^4 + 2 \cdot 3^2 x (486\theta^4 + 324\theta^3 + 279\theta^2 + 117\theta + 16) \\ & + 2^2 3^8 x^2 (972\theta^4 + 1296\theta^3 + 1035\theta^2 + 306\theta + 32) \\ & + 2^4 3^{16} x^3 (2\theta + 1)^2 (3\theta + 1) 83\theta + 2) \end{aligned}$$

$$A_n = 27^n \binom{2n}{n}^2 \binom{3n}{n} \sum_k (-1)^{n+k} \binom{n}{k} \binom{-1/3}{k} \binom{-2/3}{k} \binom{n+k}{n}^{-1}$$

155.

$$\begin{aligned} & \theta^4 + 2^4 x (3072\theta^4 + 2048\theta^3 + 1728\theta^2 + 704\theta + 87) \\ & + 2^{18} x^2 (3072\theta^4 + 4096\theta^3 + 3200\theta^2 + 896\theta + 87) \\ & + 2^{36} x^3 (2\theta + 1)^2 (4\theta + 1) (4\theta + 3) \end{aligned}$$

$$A_n = 64^n \binom{2n}{n}^2 \binom{4n}{2n} \sum_k (-1)^{n+k} \binom{n}{k} \binom{-1/4}{k} \binom{-3/4}{k} \binom{n+k}{n}^{-1}$$

165.

$$\begin{aligned} & \theta^4 - 3^2 x (33\theta^4 + 66\theta^3 + 57\theta^2 + 24\theta + 4) \\ & + 2^3 3^6 x^2 (\theta + 1)^2 (5\theta^2 + 10\theta + 4) \\ & - 2^2 3^{10} x^3 (\theta + 1) (\theta + 2) 82\theta + 1) 82\theta + 5) \end{aligned}$$

$$A_n = 27^n \binom{2n}{n} \sum_k (-1)^{n+k} \binom{n}{k} \binom{3k}{n} \binom{-1/3}{k} \binom{-2/3}{k}$$

214.

$$\begin{aligned} & \theta^4 - 2x (102\theta^4 + 204\theta^3 + 155\theta^2 + 53\theta + 7) \\ & + 2^2 x^2 (\theta + 1)^2 (396\theta^2 + 792\theta + 311) \\ & - 2^4 7^2 x^3 (\theta + 1) 8\theta + 2) 82\theta + 1) (2\theta + 5) \end{aligned}$$

$$A_n = \binom{2n}{n} \sum_{i,j} (-1)^{i+j} \binom{n}{i} \binom{n}{j} \binom{i+j}{j}^3$$

227.

$$\begin{aligned} &\theta^4 - 2^2 3^2 x(132\theta^4 + 264\theta^3 + 201\theta^2 + 69\theta + 10) \\ &+ 2^9 3^6 x^2(20\theta^4 + 80\theta^3 + 107\theta^2 + 54\theta + 10) \\ &+ 2^{12} 3^{10} x^3(2\theta + 1)^2(2\theta + 3)^2 \end{aligned}$$

$$A_n = 432^n \binom{2n}{n} \sum_k (-1)^k \binom{n}{k} \binom{3k}{n} \binom{-1/6}{k} \binom{-5/6}{k}$$

228.

$$\begin{aligned} &\theta^4 - 2^2 x(176\theta^4 + 352\theta^3 + 289\theta^2 + 113\theta + 18) \\ &+ 2^{11} x^2(80\theta^4 + 320\theta^3 + 449\theta^2 + 258\theta + 54) \\ &- 3 \cdot 2^{16} x^3(2\theta + 1)(2\theta + 5)(4\theta + 3)(4\theta + 9) \end{aligned}$$

$$A_n = 64^n \binom{2n}{n} \sum_k (-1)^{n+k} \binom{n}{k} \binom{3k}{n} \binom{-1/4}{k} \binom{-3/4}{k}$$

4. Difference equations of low order.

Most of the CY equations were found by using Maple's "Zeilberger" on a simple sum of binomial coefficients. Maple answers with a recursion formula of smallest order, but does not care if the degree is high. Since when converting to differential equations, the degree becomes the order of the differential equation, we often end up with order larger than 4 (usually 6). In most cases the differential equation L can be factored $L = L_1 L_2$ where the order of L_2 is 4.

Consider the noncommutative *algebra of difference operators*

$$\mathbf{C} \langle n, N \rangle$$

where

$$Nf(n) = f(n+1)$$

We have the relation

$$Nn = nN + N$$

We have the *Weyl algebra* of differential operators with polynomial coefficients

$$\mathbf{C} \langle x, \theta \rangle$$

where $\theta = x \frac{d}{dx}$ with the relation

$$\theta x = x\theta + x$$

Proposition.

There is an isomorphism between the difference operator algebra

$$\mathbf{C}\langle n, N \rangle / (Nn - nN - N)$$

and the Weyl algebra

$$\mathbf{C}\langle x, \theta \rangle / (\theta x - x\theta - x)$$

given by the map

$$n \rightarrow -\theta$$

$$N \rightarrow x$$

Hereby

$$order \rightarrow degree$$

$$degree \rightarrow order$$

In order to solve the differential equations we need to write them in ordinary fashion. For this the following formulas are useful

$$\theta^n = \sum_{j=1}^n S(n, j) x^j \frac{d^j}{dx^j}$$

$$x^n \frac{d^n}{dx^n} = \sum_{j=1}^n s(n, j) \theta^n$$

where $S(n, j)$ and $s(n, j)$ are *Stirling numbers* of second and first kind respectively. This gives a very nice way of defining the Stirling numbers without referring to combinatorics.

Example (#232) Let

$$A_n = \binom{2n}{n}^2 \sum_k \binom{n}{k}^2 \binom{3n}{n+k}$$

Maple gives the following difference operator

$$(n+2)^4 Q_0(n+2) N^2$$

$$-(37345n^6 + 272085n^5 + 806321n^4 + 123965n^3 + 1038758n^2 + 448272n + 77824)N$$

$$-384(2n+1)^2(3n+1)(3n+2)Q_0(n+3)$$

where

$$Q_0(n) = 77n^2 - 209n + 142$$

Converting this to a differential operator we get

$$L = \theta^4 Q_0(\theta)$$

$$\begin{aligned} & -x(37345\theta^6 + 48015\theta^5 + 6071\theta^4 - 11683\theta^3 - 2944\theta^2 + 78\theta + 240) \\ & -384x^2(2\theta + 1)^2(3\theta + 1)(3\theta + 2)Q_0(\theta + 3) \end{aligned}$$

Factorization gives

$$L = L_1 L_2$$

where

$$\begin{aligned} L_1 = & 5^2 Q_0(\theta + 4) - 5x(201509\theta^2 + 1453353\theta + 2642894) \\ & + 2^6 3x^2(51821^2 + 287078\theta + 499706) \\ & + 2^{10} 3^2 5x^3(14707\theta^2 + 158741\theta + 453284) \\ & - 2^{19} 3^5 x^4 Q_0(\theta + 7) \end{aligned}$$

and

$$\begin{aligned} L_2 = & 5^2 \theta^4 - 5x(2617\theta^4 + 4658\theta^3 + 3379\theta^2 + 1050\theta + 120) \\ & - 2^6 3x^2(-673\theta^4 + 4871\theta^3 + 10282\theta^2 + 5410\theta + 860) \\ & + 2^{10} 3^2 x^3(955\theta^4 + 4320\theta^3 + 3477\theta^2 + 1020\theta + 100) \\ & - 2^{17} 3^3 x^4(2\theta + 1)^2(3\theta + 1)(3\theta + 2) \end{aligned}$$

We also give an example of Hadamard squares. Let

$$u_0 = \sum_{n=0}^{\infty} E_n x^n$$

be the solution of

$$\theta^2 - x(a\theta^2 + a\theta + b) - cx^2(\theta + 1)^2$$

(one of the 10 cases (a),(b),...(j) in [2]). Then the Hadamard square

$$y_0 = \sum E_n^2 x^n = \sum A_n x^n$$

satisfies the recursion of order 3

$$\begin{aligned} & U(n)(n+3)^4 A_{n+3} - U(n+1) \{U(n)U(n+1) + c(n+2)^4\} A_{n+2} \\ & - cU(n) \{U(n)U(n+1) + c(n+2)^4\} A_{n+1} + c^3 U(n+1)(n+1)^4 A_n = 0 \end{aligned}$$

where

$$U(n) = a(n+1)^2 + a(n+1) + b$$

This gives a degree 3 differential equation

$$Q_0(\theta)\theta^4 - \dots + c^3 x^3 Q_0(\theta + 4)(\theta + 1)^4$$

where

$$Q_0(n) = a(n-2)^2 + a(n-2) + b$$

By duality we believe that difference operators of order 2 and 3 are worth a table which can be used as a "superseeker" to recognize CY differential equations without solving them. We order them with respect to the absolute value of the discriminant D of $Q_0(n)$. It is worth mentioning that we have found numerous formulas for the same coefficient of a CY equation. As an example we have about 40 different formulas for the coefficient A_n of #25. For hundreds of such formulas see [4].

Table. Superseeker II.

$ D $	D	$Q_0(n)$	#	degree
3	-3	$7n^2 - 23n + 19$	27	?
3	-3	$7n^2 - 19n + 13$	243	?
4	-2^2	$10n^2 - 26n + 17$	237,256	2
5	5	$5n^2 - 15n + 11$	253	2
7	-7	$14n^2 - 35n + 22$	241,33	2
12	$-2^2 3$	$12n^2 - 30n + 19$	258	2
12	$2^2 3$	$2n^2 - 2n - 1$	56	3
12	$2^2 3$	$2n^2 - 1 + n + 11$	23	3
15	$-3 \cdot 5$	$51n^2 - 147n + 106$	222	2
15	$-3 \cdot 5$	$15n^2 - 45n + 34$	216	3
15	$-3 \cdot 5$	$20n^2 - 55n + 38$	55	2
15	$-3 \cdot 5$	$24n^2 - 57n + 34$	211	2
16	-2^4	$20n^2 - 56n + 33$	119	3
20	$2^2 5$	$4n^2 - 18n + 19$	262	3
27	-3^3	$27n^2 - 63n + 37$	239	2
28	$2^2 7$	$6n^2 - 26n + 27$	235	4
32	-2^5	$24n^2 - 56n + 33$	265	2
35	$-5 \cdot 7$	$21n^2 - 49n + 29$	71	3
35	$-5 \cdot 7$	$21n^2 - 77n + 71$	21	3

$ D $	D	$Q_0(n)$	#	degree
37	37	$41n^2 - 105n + 67$	300	2
39	$-3 \cdot 13$	$20n^2 - 51n + 33$	223	3
44	$-2^2 11$	$33n^2 - 88n + 59$	278	2
44	$-2^2 11$	$44n^2 - 110n + 69$	238,288	2
55	$-5 \cdot 11$	$77n^2 - 209n + 142$	232	2
60	$-2^2 3 \cdot 5$	$40n^2 - 90n + 51$	277	2
60	$-2^2 3 \cdot 5$	$48n^2 - 126n + 83$	210	2
105	$3 \cdot 5 \cdot 7$	$12n^2 - 45n + 40$	242,259	3
135	$-3^3 5$	$27n^2 - 99n + 92$	266	3
140	$-2^2 5 \cdot 7$	$52n^2 - 134n + 87$	282	2
160	$2^5 5$	$8n^2 - 8n - 3$	83	3
176	$2^4 11$	$44n^2 - 88n + 43$	254,295	?
195	$-3 \cdot 5 \cdot 13$	$85n^2 - 235n + 163$	99	2
224	$-2^5 7$	$84n^2 - 196n + 115$	289	2
231	$-3 \cdot 7 \cdot 11$	$55n^2 - 143n + 94$	117,118	3
231	$-3 \cdot 7 \cdot 11$	$55n^2 - 187n + 160$	22,212	3
240	$-2^4 3 \cdot 5$	$204n^2 - 432n + 229$	225	2
252	$-2^2 3^2 7$	$56n^2 - 154n + 107$	215	3
255	$-3 \cdot 5 \cdot 17$	$87n^2 - 327n + 308$	279	3

$ D $	D	$Q_0(n)$	#	degree
288	$2^5 3^2$	$8n^2 - 40n + 41$	119	3
320	$-2^6 5$	$48n^2 - 176n + 163$	246	3
320	$-2^6 5$	$48n^2 - 112n + 67$	247	3
345	$3 \cdot 5 \cdot 23$	$92n^2 - 299n + 242$	226	3
385	$5 \cdot 7 \cdot 11$	$44n^2 - 143n + 114$	219	3
399	$-3 \cdot 7 \cdot 19$	$285n^2 - 969n + 824$	59	3
399	$-3 \cdot 7 \cdot 19$	$102n^2 - 309n + 235$	218	3
455	$-5 \cdot 7 \cdot 13$	$156n^2 - 403n + 261$	109	2
495	$-3^2 5 \cdot 11$	$170n^2 - 415n + 254$	192	2
495	$-3^2 5 \cdot 11$	$88n^2 - 231n + 153$	260	3
640	$2^7 5$	$32n^2 - 64n + 27$	261	3
1463	$-7 \cdot 11 \cdot 19$	$171n^2 - 551n + 446$	198	?
1564	$-2^2 17 \cdot 23$	$184n^2 - 414n + 235$	264	2
1664	$-2^7 13$	$364n^2 - 780n + 419$	294	2
2156	$2^2 7^2 11$	$110n^2 - 370n + 313$	217	3
2176	$2^7 17$	$96n^2 - 176n + 75$	276	2
2560	$-2^9 5$	$128n^2 - 416n + 343$	275	3
2665	$5 \cdot 13 \cdot 41$	$164n^2 - 533n + 429$	274	3
3135	$-3 \cdot 5 \cdot 11 \cdot 19$	$154n^2 - 407n + 274$	231	3

$ D $	D	$Q_0(n)$	#	degree
3335	$-5 \cdot 23 \cdot 29$	$203n^2 - 551n + 378$	224	3
4180	$2^2 5 \cdot 11 \cdot 19$	$231n^2 - 484n + 249$	230	3
5831	$-7^3 17$	$340n^2 - 1037n + 795$	234	3
9204	$2^2 3 \cdot 13 \cdot 59$	$295n^2 - 944n + 763$	248	3
14400	$-2^6 3^2 5^2$	$288n^2 - 936n + 773$	249	3
17199	$-3^3 7^2 13$	$572n^2 - 1599n + 1125$	297	3
17415	$-3^4 5 \cdot 43$	$324n^2 - 1053n + 869$	273	3
39767	$-7 \cdot 13 \cdot 19 \cdot 23$	$884n^2 - 2405n + 1647$	208	3
44591	$-17 \cdot 43 \cdot 61$	$946n^2 - 2623n + 1830$	209	3
60480	$2^6 3^3 5 \cdot 7$	$432n^2 - 1584n + 1487$	268	3
64496	$-2^4 29 \cdot 139$	$1740n^2 - 3712n + 1989$	305	2
64844	$-2^2 13 \cdot 29 \cdot 43$	$812n^2 - 2378n + 1761$	240	2
104719	$-23 \cdot 29 \cdot 157$	$1595n^2 - 5191n + 4240$	19	3
170375	$5^3 29 \cdot 47$	$1457n^2 - 4277n + 3168$	195	?
702075	$-3 \cdot 5^2 \cdot 11 \cdot 23 \cdot 37$	$1771n^2 - 4807n + 3361$	252	3
959040	$-2^6 3^4 5 \cdot 37$	$2592n^2 - 8424n + 6937$	272	3
5274751	$23 \cdot 79 \cdot 2903$	$6557n^2 - 18565n + 13342$	250	3

Acknowledgements.

Firstly I want to thank Christian van Enckevort and Duco van Straten who found many of the equations listed here. Further I thank Wadim Zudilin who has inspired much of this work and communicated the pullback of Yinfan to me. Finally I thank Yinfan Yang , whose pullback caused the writing of this paper.

References.:

1. G.Almkvist, Strängar i månsken I, Normat 51 (2003), 22-33, II, Normat 51 (2003), 63-79
2. G.Almkvist, W.Zudilin, Differential equations, mirror maps and zeta values. In Mirror symmetry V, Proceedings of BIRS workshop on Calabi-Yau Varieties and Mirror Symmetry, December 6-11, 2003
3. G.Almkvist,C. van Enckevort, D. van Straten, W.Zudilin, Tables of Calabi-Yau equations, math.AG/0507430
4. G.Almkvist, Some binomial identities related to Calabi-Yau differential equations, In preparation
5. C.van Enckevort, D.van Straten, Monodromy calculations for fourth order equations of Calabi-Yau type. In Mirror Symmetry V, Proceedings of BIRS workshop on Calabi-Yau Varieties and Mirror Symmetry, December 6-11, 2003.
6. Yinfan Yang, Pullbacks, personal communication
7. D.Zagier, Integral solutions of Apéry-like recurrence equations, Manuscript 2003

Math Dept
Univ of Lund
Box 118
22100 Lund, Sweden
gert@maths.lth.se